



AVISCERA BIOSCIENCE

Anti Vasostatin-2/Chromogranin A (19-131) Purified IgG

Product Information

Code	A00084-01-100
Name	hCHGA(19-131) Antibody
Clone No.	Polyclonal
Lot No.	
Size	100 µg
Species	Human
Host	Rabbit
Immunogen	hCHGA(19-131) (H), rec.
Ab Type	Purified IgG
Purification	Protein A
Formulation	Lyophilized form without preservatives free
Carry	
Storage	-20 ° C
Specificity	H, R, M
Reconstitution	PBS, 100 µl
Application	IHC WB ELISA

AVISCERA BIOSCIENCE INC.
2348 Walsh Ave. Suite C
Santa Clara, CA 95051
Tel: (408) 982 0300
Fax: (408) 982 0301
Email:
Sales@AvisceraBioscience.com
www.AvisceraBioscience.com

Preparation

This antibody was produced from a rabbit immunized with purified, *E. coli*-derived, recombinant human Vasostatin-2 /Chromogranin A (19-131). That antibody was purified by Protein A affinity.

Formulation

100 µg IgG in 100 µl of PBS without preservatives was lyophilized.

Reconstitution

Add 100 µl of PBS to the vial to prepare antibody stock solution at 100 µg/ µl. Store reconstituted antibody at 2 to 8 ° C for up a few weeks. This antibody can also be aliquotted (by 10 µL per vial) and stored frozen at -20° C to -70° C **in a manual defrost freezer** for up six months without detectable loss of activity.

Storage

Lyophilized antibody can be stored at 2 ~8 ° C for a few weeks or at -20 ° C for six months. **Avoid repeated freeze-thaw cycles.**

Specificity

This antibody has been selected for its ability to recognize human Vasostatin-2/Chromogranin A (19-131) in direct ELISAs.

Applications

Direct ELISA - This antibody can be used at 1:4000 (0.25 µg/ml) with the appropriate secondary reagents to detect recombinant human Vasostatin-2.

Western Blot - This antibody can be used at 1:500~1000 (1~2 µg/ml) with the appropriate secondary reagents to detect recombinant human Vasostatin-2 under reduce conditions. The detection limit for rh Vasostatin-2 approximately 15 ng/lane under reducing conditions.

Immunohistochemistry - This antibody can be used at 1:500 (2 µg/ml) with the appropriate secondary antibody to detect Vasostatin-2 in human pancreas tissues (ABC).

Optimal dilutions should be determined by each laboratory for each application.