

#### **Product Information**

A00766-16-

100

Name Human SPARC

Mab

Clone No. 77-9

Lot No.

Size 100 μg

Species Human

Host Mouse

Immunogen Human SPARC,

rec.

Ab Type IgG

Purification sequential

precipitation Lyophilized

Formulation Form without

preservatives

Carry Free

Storage -20 ° C

Specificity Human

Reconstitution 100  $\mu$ l

Application ELISA

AVISCERA BIOSCIENCE INC. 2348 Walsh Ave. Suite C Santa Clara, CA 95051 Tel: (408) 982 0300 Fax: (408) 982 0301

Email:

Sales@AvisceraBioscience.com www.AvisceraBioscience.com

# **AVISCERA BIOSCIENCE**

# Anti Human Secreted Protein Acidic and Rich in Cysteine (SPARC)/osteonectin Monoclonal IgG

# **Preparation**

This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, E. coli-derived, recombinant human SPARC, His Tag on N-Terminal. That antibody was purified by sequential precipitation caprylic acid and ammonium sulphate.

#### **Formulation**

100 μg of mouse IgG in 100 μl of PBS lyophilized form.

# **Reconstitution and Storage**

Add 100  $\mu$ l deionized water to the vial to prepare a antibody stocking solution (100 $\mu$ g/ml). Stores it at 4°C for a few days. For long term storage, the reconstituted antibody can also be aliquotted ( by 10  $\mu$ L per vial) and stored frozen at -20° C to -70° C in a manual defrost freezer for 12 months without detectable loss of activity. Avoid repeated freeze-thaw cycles.

## Specificity

This antibody has been selected for its ability to recognize recombinant human SPARC in indirect ELISAs.

### **Applications**

**Indirect ELISA** - This antibody can be used at 1:2187000 (0.45 ng/ml) to detect human SPARC on indirectly ELISA.

**ELISA Assay** - This antibody can be used as a capture antibody in a human SPARC sandwich immunoassay in combination with the human SPARC detection antibody (Code No.: A00766-12-50B) and recombinant human SPARC (Code No.: 00766-01-100) as the standard. The suggested concentration range for this capture antibody is 0.5 μg/mL and should be titrated to determine the optimal concentration.

Optimal dilutions should be determined by each laboratory for each application.

THIS PRODUCT IS FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.