



AVISCERA BIOSCIENCE

Rabbit Anti Betatrophin Isoform CRA_a (Human) Purified IgG

Product Information

Code	A00528-17-100
Name	Human Betatrophin Isoform CRA_a Antibody
Clone No.	Polyclonal
Lot No.	
Size	100 µg
Species	Human
Host	Rabbit
Immunogen	Human Betatrophin Isoform CRA_a.
Ab Type	Purified IgG
Purification	Protein A
Formulation	Lyophilized form without preservatives free
Carry	
Storage	-20 ° C
Specificity	Human
Reconstitution	PBS, 100 µl
Application	IHC WB ELISA

AVISCERA BIOSCIENCE INC.
2348 Walsh Ave. Suite C
Santa Clara, CA 95051
USA
Tel: (408) 982 0300
Fax: (408) 982 0301
Email:
Sales@AvisceraBioscience.com
www.AvisceraBioscience.com
www.AvisceraBioscience.net

Preparation

This antibody was produced from a rabbit immunized with purified recombinant human Betatrophin Isoform CRA_a. That antibody was purified by Protein A affinity.

Formulation

100 µg of Rabbit Anti Human Betatrophin Isoform CRA_a IgG in 100 µl of PBS without preservatives was lyophilized.

Reconstitution

Add 100 µl of PBS to the vial to prepare antibody stock solution at 100 µg/ 100µl. Store reconstituted antibody at 2 to 8 ° C for up a few weeks. This antibody can also be aliquotted (by 10 µL per vial) and stored frozen at -20° C to -70° C in a **manual defrost freezer** for up 12 months without detectable loss of activity.

Storage

Lyophilized antibody can be stored at 2 ~8 ° C for a few weeks or at -20 ° C for six months. **Avoid repeated freeze-thaw cycles.**

Specificity

This antibody has been selected for its ability to recognize human Betatrophin Isoform CRA_a on indirect ELISA, Immunohistochemistry and western blot.

Applications

Indirect ELISA - This antibody can be used at 0.5 µg/ml with the appropriate secondary reagents to detect recombinant human Betatrophin Isoform CRA_a.

Western Blot - This antibody can be used at 4-6 µg/ml with the appropriate secondary reagents to detect recombinant human Betatrophin Isoform CRA_a under reduce conditions. The detection limit for rh Betatrophin Isoform CRA_a approximately 28 ng/lane under reducing conditions.

Immunohistochemistry - This antibody can be used at 4-6 µg/ml with the appropriate secondary antibody to detect Betatrophin Isoform CRA_a in the paraffin embedded human hepatocellular carcinoma tissues and visceral adipose tissues (ABC).

THIS PRODUCT IS FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.