



## Rabbit Anti-CTRP13 (Human, Mouse) IgG

### Product Information

Code	A00333-01-100
Name	Anti CTRP13 (H,M) IgG
Clone No.	N/A
Lot No.	
Size	100 µg
Species	Human, Mouse
Host	Rabbit
Immunogen	Human CTRP13 rec.
Ab Type	IgG
Purification	Protein A
Formulation	Lyophilized Form without Preservatives
Carry	Free
Storage	-20 ° C
Specificity	H, R, M
Reconstitution	PBS, 100 µl
Application	IHC  ELISA

AVISCERA BIOSCIENCE INC.  
2348 Walsh Ave. Suite C  
Santa Clara, CA 95051  
Tel: (408) 982 0300  
Fax: (408) 982 0301  
Email:  
Sales@AvisceraBioscience.com  
www.AvisceraBioscience.com

### Preparation

This antibody was produced from a rabbit immunized with purified recombinant human CTRP13. That IgG was purified by Protein A affinity.

### Formulation

100µg of purified Anti CTRP13 (Human, Mouse) IgG in 100 µl of PBS without preservatives was lyophilized.

### Reconstitution

Add 100 µl of PBS to the vial to prepare antibody stock solution at 100 µg/100 µl. Store reconstituted antibody at 2 to 8 ° C for up a few weeks. This antibody can also be aliquotted ( by 10 µL per vial) and stored frozen at -20° C to -70° C in a **manual defrost freezer** for up six months without detectable loss of activity.

### Storage

Lyophilized antibody can be stored at 2 ~8 ° C for a few weeks or at -20 ° C for six months. **Avoid repeated freeze-thaw cycles.**

### Specificity

This antibody has been selected for its ability to recognize CTRP13 (Human, Mouse) on indirect ELISA, Immunohistochemistry.

### Applications

**Indirect ELISA** - This antibody can be used at 0.125 µg/mL with the appropriate secondary reagents to detect CTRP13 (Human, Mouse) on indirect ELISA.

**Immunohistochemistry**-That Antibody can be used at 2- 4 µg /mL with the appropriate secondary antibody to detect CTRP13 in the paraffin embedded human, rat or mouse adipose tissues (ABC).

**ELISA**-This antibody can be used as a detection antibody combines with the capture antibody A00333-04-100 or A00333-05-100 to detect CTRP13 (Human, Mouse) on ELISA.

*Optimal dilutions should be determined by each laboratory for each application.*